

Estimation of qubit pure states with collective and individual measurements

E. Bagan, A. Monras and R. Muñoz-Tapia

*Grup de Física Teòrica & IFAE, Facultat de Ciències, Edifici Cn,
Universitat Autònoma de Barcelona, 08193 Bellaterra (Barcelona) Spain*

We analyze the estimation of a qubit pure state by means of local measurements on N identical copies and compare its average fidelity for an isotropic prior probability distribution to the absolute upper bound given by collective measurements. We discuss two situations: the first one, where the state is restricted to lie on the equator of the Bloch sphere, is formally equivalent to phase estimation; the second one, where there is no constrain on the state, can also be regarded as the estimation of a direction in space using a quantum arrow made out of N parallel spins. We discuss various schemes with and without classical communication and compare their efficiency. We show that the fidelity of the most general collective measurement can always be achieved asymptotically with local measurements and no classical communication.

PACS numbers: 03.67.-a, 03.65.Wj, 89.70.+c

I. INTRODUCTION

In Quantum Information, measurement and estimation are not just *some* important topics. They are at the very core of the theory. An unknown quantum state can only be unveiled by means of measurements, and the information is always gained at the expense of destroying the state. This information is then processed to obtain the desired estimate.

Any estimation procedure requires a sample of identical copies of the unknown state on which we can perform measurements. If unknown states could be copied, then one could produce samples of an arbitrary number of copies and the state could be estimated with infinite accuracy. The no-cloning theorem, however, prevents this possibility [1]. But even in such unphysical circumstances, one would only have finite time and a limited number of resources for copying and measuring. So, in the real world only a finite, usually not large, number of copies are available and only a reasonable approximation to the unknown state can be made. It is thus very important to devise strategies with optimal performance in a variety of practical situations.

Over the last few years, it has been recognized that a joint measurement on N copies is more efficient than N individual measurements on each copy separately. We will often refer to the former as *collective*, in contrast to *individual* (also called *local*) measurements. The quantum correlations behind the collective measurements are (almost) always more powerful than the classical correlations used in sequential individual measurements [2, 3]. This and other issues have been studied in various contexts [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17], but for local measurements not many analytical results have been obtained [18, 19, 20, 21, 22, 23, 24]. The most powerful, involve sophisticated estimation theory technology that is not so widely known among physicists. Moreover, they mainly apply in the asymptotic limit (when N is large), and in a “pointwise” fashion, in the sense that no average over the prior probability distribution of unknown states is considered [25, 26].

Here we address the issue of estimating the most elementary quantum pure state, that of a qubit, assuming we have a sample of N identical copies of it. We consider two relevant cases: estimation of a completely unknown qubit state, i.e. one given by an arbitrary point on the Bloch sphere; and estimation of restricted states that are known to lie on the equator of the Bloch sphere. The latter is also interesting because it is formally equivalent to phase estimation (these states are also equivalent to the so-called rebits [27]). We refer to these two situations as 3D and 2D cases, respectively.

The most general measurement is described by a Positive Operator Valued Measure (POVM) on the N copies. An optimal measurement of this type yields the ultimate bounds that can be achieved by any estimation procedure. We will re-derive these bounds in a unified, very comprehensive framework. For local measurements, we will consider von Neumann measurements, as these can be readily implemented in a laboratory. So, by “local measurement” we will loosely mean a *local von Neumann measurement*. Furthermore, we will show that for some of the local procedures discussed here, optimal measurements are necessarily of von Neumann type. We will refer to local schemes that use classical communication, i.e. those that exploit the possibility of using the actual outcomes that are being obtained to dynamically adapt the next measurements, as LOCC (Local Operations and Classical Communication) schemes.

Let us be more concrete about the problem we are concerned with. Assume that we are given an ensemble of N identical copies of an unknown qubit state, which we denote by $|\vec{n}\rangle$, where $\vec{n}$ is the unit vector on the Bloch sphere that satisfies

$$|\vec{n}\rangle\langle\vec{n}| = \frac{1 + \vec{n} \cdot \vec{\sigma}}{2}, \quad (1)$$

and $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ are the usual Pauli matrices. After performing a measurement (collective or local) on the N copies of $|\vec{n}\rangle$, one obtains an outcome χ . Based on χ , an estimate, $|\vec{M}(\chi)\rangle$, for the unknown state is guessed. To quantify how well $|\vec{M}(\chi)\rangle$ approximates the unknown

state $|\vec{n}\rangle$ we use the fidelity, defined as the overlap

$$f_n(\chi) \equiv |\langle \vec{n} | \vec{M}(\chi) \rangle|^2 = \frac{1 + \vec{n} \cdot \vec{M}(\chi)}{2}. \quad (2)$$

The fidelity (2) can be regarded as a “score”: we get “1” for a perfect determination ($\vec{M} = \vec{n}$) and “0” for a completely wrong guess ($\vec{M} = -\vec{n}$). Our aim is to maximize the average fidelity, hereafter fidelity in short, over the initial probability and all possible outcomes,

$$F \equiv \langle f \rangle = \sum_{\chi} \int dn f_n(\chi) p_n(\chi), \quad (3)$$

where dn is the prior probability distribution, and $p_n(\chi)$ is the probability of getting outcome χ given that the unknown state is $|\vec{n}\rangle$.

As mentioned above, we allow for general measurements or POVMs. They are defined by a set of positive operators $\{O(\chi)\}$ (each one of them associated to an outcome) that satisfy the condition

$$\sum_{\chi} O(\chi) = \mathbb{1}. \quad (4)$$

The probability $p_n(\chi)$ is given in terms of these operators by

$$p_n(\chi) = \text{tr} [\rho_n O(\chi)], \quad (5)$$

where ρ_n is the quantum state of the N identical copies, i.e., $\rho_n = (|\vec{n}\rangle\langle\vec{n}|)^{\otimes N}$.

In Eq. (3) there are only two elements that require optimization: the guess and the POVM, which enter this equation through (2) and (5), respectively. The optimal guess (OG) can be obtained rather trivially. The Schwarz inequality shows that the choice

$$\vec{M}(\chi) = \frac{\vec{V}(\chi)}{|\vec{V}(\chi)|}, \quad (6)$$

where

$$\vec{V}(\chi) = \int dn \vec{n} p_n(\chi), \quad (7)$$

maximizes the value of the fidelity, which then reads [22]

$$F = \frac{1}{2} (1 + \Delta) \equiv \frac{1}{2} \left(1 + \sum_{\chi} |\vec{V}(\chi)| \right) \quad (8)$$

(in the strict sense we should write F_{OG} , but to simplify the notation, we drop the subscript when no confusion arises). Eq. (6) gives the best state that can be inferred, and Eq. (8) gives the maximum fidelity that can be achieved for *any* prior probability and *any* measurement scheme specified by the conditional probabilities $p_n(\chi)$. We are thus left only with the non-trivial task of obtaining the optimal measurement. The goal of this

paper is to compute the maximum value of (8) within a unified framework for various measurement schemes, specially the local ones.

The paper is organised as follows. In the next section we derive the bounds on the fidelity of the optimal collective measurements. In Sec. III we discuss several local measurement schemes, with and without classical communication. Asymptotic values of the fidelity are computed in Sec. IV. A summary of results and our main conclusions are presented in Sec. V, and two technical appendices end the paper.

II. COLLECTIVE MEASUREMENTS

A. 2D states

A 2D state corresponds to a point that is known to lay on the equator of the Bloch sphere. If we take it to be on the xy plane, such state, $|\vec{n}\rangle$, has $\vec{n} = (\cos \theta, \sin \theta, 0)$. If no other information is available, the prior probability distribution has to be isotropic, that is, $dn = d\theta/(2\pi)$. Eq. (8) then reads

$$\Delta = \sum_{\chi} \left| \int \frac{d\theta}{2\pi} \vec{n} \text{tr} [\rho_n O(\chi)] \right|. \quad (9)$$

Notice that we can write,

$$\rho_n \equiv \rho(\theta) = U(\theta) \rho_0 U^\dagger(\theta), \quad (10)$$

where ρ_0 is a fiducial state of angular momentum $J \equiv N/2$ and maximal magnetic quantum number, $m = J$, along any fixed direction on the equator of the Bloch sphere. In particular, ρ_0 can be chosen to point along the x axis,

$$\rho_0 = |JJ\rangle_{xx} \langle JJ|. \quad (11)$$

In Eq. (10), $U(\theta)$ is the unitary representation of a rotation around the z axis. The group of such unitary matrices is isomorphic to $U(1)$. In the standard basis $|jm\rangle \equiv |m\rangle$, where $U(\theta)$ is diagonal, we have

$$\rho(\theta) = \sum_{m,n} e^{i(m-n)\theta} (\rho_0)_{mn} |m\rangle\langle n|. \quad (12)$$

We are now ready to compute (9):

$$\begin{aligned} \Delta &= \sum_{\chi} \left| \sum_{mn} \int \frac{d\theta}{2\pi} e^{i\theta} e^{i(m-n)\theta} (\rho_0)_{mn} [O(\chi)]_{nm} \right| \\ &= \sum_{\chi} \left| \sum_{m=-J}^{J-1} (\rho_0)_{mm+1} [O(\chi)]_{m+1,m} \right|, \end{aligned} \quad (13)$$

where $[O(\chi)]_{mn} \equiv \langle m | O(\chi) | n \rangle$. The following inequalities give the maximum value of Δ :

$$\Delta \leq \sum_{\chi} \sum_{m=-J}^{J-1} |(\rho_0)_{mm+1} [O(\chi)]_{m+1,m}|$$

$$\begin{aligned}
&\leq \sum_{m=-J}^{J-1} |(\rho_0)_{mm+1}| \sum_{\chi} |[O(\chi)]_{m+1m}| \\
&\leq \sum_{m=-J}^{J-1} |(\rho_0)_{mm+1}|, \tag{14}
\end{aligned}$$

where in the last step in (14) we have used that

$$\sum_{\chi} |[O(\chi)]_{m+1m}| \leq 1, \tag{15}$$

as follows from positivity and (4). More precisely, positivity implies

$$[O(\chi)]_{mm}[O(\chi)]_{m+1m+1} \geq |[O(\chi)]_{mm+1}|^2, \tag{16}$$

and the Schwarz inequality yields

$$\begin{aligned}
&\sum_{\chi} |[O(\chi)]_{m+1m}| \leq \\
&\sqrt{\sum_{\chi} [O(\chi)]_{mm}} \sqrt{\sum_{\chi} [O(\chi)]_{m+1m+1}} = 1. \tag{17}
\end{aligned}$$

There are two points worth emphasizing here. First, all the inequalities can be saturated, therefore the bound is tight (we give below a POVM that accomplishes this task). Second, the bound (14) is completely general. If we were interested in encoding the information carried by a phase θ in a covariant way [as in (10)], but using a general fiducial state [7, 28, 29], $\rho_0 = \sum_{mm'} a_m a_{m'}^* |m\rangle\langle m'|$ (ideally the best possible one, which is not necessarily a tensor product of identical copies), the fidelity would still be bounded by (14). In this general case, one has $\Delta \leq \sum_{mm'} a_{m'}^* \mathbf{M}_{m'm} a_m$, where $2\mathbf{M}_{m'm} = \delta_{m'm+1} + \delta_{m'+1m}$, and its maximum value is given by the largest eigenvalue of the matrix $\mathbf{M}$. A straightforward calculation gives [28] $F_{\max} = (1 + \cos[2\pi/(d_J + 2)])/2$, where $d_j = 2j + 1$ is the dimension of the invariant Hilbert space corresponding to the representation $\mathbf{j}$ of $SU(2)$.

In our problem ρ_0 is constrained by the condition of having identical copies. From (11) and (14) one has [8]

$$\begin{aligned}
\Delta &\leq \frac{1}{2^N} \sum_{m=-J}^{J-1} \sqrt{\binom{N}{J+m} \binom{N}{J+m+1}} \\
&= \frac{1}{2^N} \sum_{m=-J}^J \binom{N}{J+m} \sqrt{\frac{J-m}{J+m+1}}, \tag{18}
\end{aligned}$$

where we have used that

$$|JJ\rangle_x = |\vec{x}\rangle^{\otimes N} = \frac{1}{2^J} \sum_{m=-J}^J \binom{N}{J+m} |Jm\rangle \tag{19}$$

(recall that $J \equiv N/2$).

We next show that there are POVMs that attain this bound. To saturate the first inequality in (14) the phase

of $[O(\chi)]_{m+1m}$ must be independent of m . This is ensured if this phase is a function of $m - n$. Similarly, a set of positive operators for which $|O(\chi)_{mn}| = \text{constant}$ for all χ, m and n , will certainly saturate the remaining inequalities in (14). In particular, the covariant (continuous) POVM, whose elements are given by

$$[O(\phi)]_{mn} = e^{i(m-n)\phi}, \tag{20}$$

satisfies all the requirements. Note that we have labeled the outcomes by a rotation angle ϕ , which plays the role of χ . Hence, condition (4) becomes

$$\int \frac{d\phi}{2\pi} O(\phi) = \mathbb{1}, \tag{21}$$

which certainly holds for (20). These are rank one operators, and can also be written as

$$O(\phi) = U(\phi)|B\rangle\langle B|U^\dagger(\phi), \tag{22}$$

where

$$|B\rangle = \sum_{m=-J}^J |J, m\rangle. \tag{23}$$

We have just shown that at least one optimal measurement [i.e., a POVM that saturates (14)] exists, but other optimal measurements can be found. POVMs with finite number of outcomes, for instance, are straightforward to obtain by choosing ϕ to be the d_J -th roots of unity, namely,

$$[O(k)]_{mn} = \frac{1}{d_J} \exp\left\{i(m-n)\frac{2\pi k}{d_J}\right\}, \tag{24}$$

where $k = 1, \dots, d_J$. In this case $\{O(k)\}$ is a von Neumann measurement, since the number of rank one POVM elements equals the dimension of the Hilbert space.

In the asymptotic limit the fidelity can be obtained in terms of the moments of a binomial distribution $\text{Bin}(n, p)$ with parameters $n = N$ and $p = 1/2$. We simply need to expand (18) in powers of m , i.e., around $\langle m \rangle = 0$, to obtain

$$\begin{aligned}
\Delta &\leq \frac{1}{2^N} \sum_m \binom{N}{J+m} \times \\
&\left[1 - \frac{2m}{N} + \left(\frac{2m^2}{N^2} - \frac{1}{N}\right) + O(1/N^{3/2})\right] \tag{25}
\end{aligned}$$

(notice that sum over m is shifted by J with respect to the usual binomial distribution). The moments are well known to be $\langle 1 \rangle = 1$, $\langle m \rangle = 0$ and $\langle m^2 \rangle = N/4$. The latter shows that m has “dimensions” of $\sqrt{N}$, which helps to organise the expansion in powers of $1/N$. We finally obtain

$$\Delta_{\max} = 1 - \frac{1}{2N} + \dots, \tag{26}$$

and a fidelity

$$F = 1 - \frac{1}{4N} + \dots. \tag{27}$$

B. 3D states

A 3D state $|\vec{n}\rangle$ corresponds to a general point on the Bloch sphere. As in the previous section we write

$$\rho_n \equiv \rho(\vec{n}) = U(\vec{n})\rho_0 U^\dagger(\vec{n}), \quad (28)$$

where for convenience ρ_0 is now chosen to point along the z axis, i.e.,

$$\rho_0 = |JJ\rangle\langle JJ|, \quad (29)$$

and $U(\vec{n})$ is the unitary representation [i.e., the element of $SU(2)$] of the rotation that brings $\vec{z}$ into $\vec{n}$ (a rotation around the vector $\vec{z} \times \vec{n}$).

Recalling (5) and (7), we have

$$\vec{V}(\chi) = \int dn \vec{n} \text{tr} [\rho_n O(\chi)], \quad (30)$$

where dn is the invariant measure on the two-sphere, e.g.,

$$dn = \frac{d(\cos \theta) d\phi}{4\pi}, \quad (31)$$

where θ and ϕ are the standard azimuthal and polar angles. Notice that we can always define an operator $\Omega(\chi)$ in such a way that

$$O(\chi) = U[\vec{M}(\chi)]\Omega(\chi)U^\dagger[\vec{M}(\chi)], \quad (32)$$

where $\vec{M}(\chi)$ is given by (6). Taking into account that Δ is rotationally invariant one obtains

$$\Delta = \sum_\chi \left| \int dn n_z \text{tr} [\rho_n \Omega(\chi)] \right|. \quad (33)$$

We readily see that $n_z = \cos \theta = \mathfrak{D}_{00}^{(1)}(\vec{n})$, where the rotation matrices $\mathfrak{D}_{mm'}^{(j)}$ are defined in the standard way, $\mathfrak{D}_{mm'}^{(j)}(\vec{n}) = \langle jm|U(\vec{n})|jm'\rangle$. We then have

$$\begin{aligned} \Delta &= \sum_\chi \left| \sum_{mm'} \int dn \mathfrak{D}_{00}^{(1)}(\vec{n}) (\rho_n)_{mm'} \Omega_{m'm}(\chi) \right| \\ &= \sum_m \left[\sum_\chi \Omega_{mm}(\chi) \right] \int dn \mathfrak{D}_{00}^{(1)}(\vec{n}) (\rho_n)_{mm}, \end{aligned} \quad (34)$$

where in the second equality we have used that

$$\int dn \mathfrak{D}_{00}^{(1)}(\vec{n}) (\rho_n)_{mm'} = \delta_{mm'} \int dn \mathfrak{D}_{00}^{(1)}(\vec{n}) (\rho_n)_{mm}, \quad (35)$$

as follows from Schur's lemma after realizing that the left hand side of (35) commutes with $U(\theta)$, the unitary transformations defined right after (11). Recall that these transformations, which are a $U(1)$ subgroup of $SU(2)$, have only one-dimensional irreducible representations, labeled by the magnetic quantum number m , thus yielding relation (35). In (34) we have removed the absolute value

as all terms are positive (see below). Tracing (32) one obtains $\sum_\chi \text{tr} \Omega(\chi) = \sum_\chi \text{tr} O(\chi) = d_J$. Therefore

$$\begin{aligned} \Delta &\leq d_J \max_m \int dn \mathfrak{D}_{00}^{(1)}(\vec{n}) (\rho_n)_{mm} \\ &= \max_m \langle 10; Jm | Jm \rangle^2 = \frac{J}{J+1}, \end{aligned} \quad (36)$$

where in the second equality we have used that $\rho_{mm} = \mathfrak{D}_{mJ}^{(J)}(\vec{n})\mathfrak{D}_{mJ}^{(J)*}(\vec{n})$ and the well known orthogonality relations of the $SU(2)$ irreducible representations [34]. Recalling that $J \equiv N/2$ we finally get [3]

$$F = \frac{N+1}{N+2}, \quad (37)$$

which for large N behaves as

$$F = 1 - \frac{1}{N} + \dots \quad (38)$$

Let us finally give a POVM that saturates all the inequalities. The maximum of the Clebsch-Gordan $\langle 10; Jm | Jm \rangle$ in (36) occurs at $m = J$. Hence, to attain the bound we need to choose

$$\Omega_{mm}(\chi) = c_\chi \delta_{Jm}, \quad \sum_\chi c_\chi = d_J, \quad (39)$$

where the coefficients c_χ are positive. This leads straightforwardly to the optimal continuous POVM defined by

$$O(\vec{m}) = d_J U(\vec{m}) |JJ\rangle\langle JJ| U^\dagger(\vec{m}) \quad (40)$$

(one can check that $\int dm O(\vec{m}) = \mathbb{1}$). POVM's with finite number of elements can also be constructed. The only requirements are (39) and, of course, (4). These constraints translate into a series of conditions for the set of directions $\{\vec{m}\}$, for which solutions with a finite number of elements can be found. We refer the reader to the literature [8, 9, 11, 15] for details.

III. LOCAL MEASUREMENTS

Collective measurements, although very interesting from the theoretical point of view, are difficult to implement in practice. Far more interesting for experimentalists are individual von Neumann measurements [21, 31].

Individual von Neumann measurements on qubits are represented by two projectors

$$O(\pm\vec{m}) = \frac{1 \pm \vec{m} \cdot \vec{\sigma}}{2}, \quad (41)$$

where $\vec{m}$ is a unit Bloch vector characterizing the measurement (in a spin system, e.g., $\vec{m}$ is the orientation of a Stern-Gerlach). In a general frame we must also allow classical communication, i.e., the possibility of adapting the orientation of the measuring devices depending on previous outcomes [20, 21, 22].

In the next sections we study quantitatively several schemes in ascending order of optimality: from the most basic tomography inspired schemes [30, 33] to the most general individual measurement procedure with classical communication [22].

Our aim is to investigate how good these local measurements are as compared to the collective ones. We would like to stress that in this context few analytical results are known [19, 23, 26]. Our results here complement and extend the analysis carried out by some of the authors in [22].

A. Fixed Measurements

Let us start with the most basic scheme for reconstructing a qubit: fixed von Neumann measurements along 2 orthogonal directions (say, x and y) in the equator of the Bloch sphere for 2D states, or along 3 orthogonal directions (say, x , y and z) for 3D states. This kind of scheme is often called tomography [30, 31, 32].

Consider $N = 2N$ ($3N$) copies of the state $|\vec{n}\rangle$. After N measurements along each one of the directions $\vec{e}_i$, $i = x, y, (z)$, we obtain a set of outcomes $+1$ and -1 with frequencies $N\alpha_i$ and $N(1 - \alpha_i)$, respectively. This occurs with probability

$$p_n(\alpha) = \prod_{i=x,y,(z)} \binom{N}{N\alpha_i} \left(\frac{1+n_i}{2}\right)^{N\alpha_i} \left(\frac{1-n_i}{2}\right)^{N(1-\alpha_i)}, \quad (42)$$

where n_i are the projections of the vector $\vec{n}$ along each direction, $n_i \equiv \vec{n} \cdot \vec{e}_i$, and we have used the shorthand notation $\alpha = \{\alpha_i\}$. The combinatorial factor takes into account all the possible orderings of the outcomes and the remaining factors are the quantum probabilities, i.e., the appropriate powers of $\text{tr} [|\vec{n}\rangle\langle\vec{n}|O(\pm\vec{e}_i)]$.

Since the expectation value of $\vec{\sigma}$ is $\langle\vec{n}|\vec{\sigma}|\vec{n}\rangle = \vec{n}$, a straightforward guess based on the relative frequencies of each outcome is

$$M_i^{\text{CLG}}(\alpha) = \frac{2\alpha_i - 1}{\sqrt{\sum_j (2\alpha_j - 1)^2}}, \quad (43)$$

where the superscript stands for central limit guess (CLG). Notice the normalization factor which ensures that $|\vec{M}^{\text{CLG}}| = 1$, hence $\vec{M}^{\text{CLG}}$ always corresponds to a physical pure state [33]. The average fidelity in this case is given by

$$F = \frac{1}{2} + \frac{1}{2} \sum_{\alpha} \int d\vec{n} \cdot \vec{M}^{\text{CLG}}(\alpha) p_n(\alpha), \quad (44)$$

where $p_n(\alpha)$ is defined in (42). Although the CLG (43) is not the *best* state that one can infer from the data, it has the nice property that it can be directly (and easily) obtained from the observed frequencies without further processing. So, it is interesting to compare its fidelity with the optimal one, given by (7) and (8) (see Fig. 1).




FIG. 1: Average fidelities in terms of the number of copies in the 2D case for the optimal collective measurement (solid line), tomographic OG (dot-dashed line), tomographic CLG (dashed line) and a simulation of the greedy scheme (dots).

The OG for this measurement scheme is $\vec{M}^{\text{OG}}(\alpha) = \vec{V}(\alpha)/|\vec{V}(\alpha)|$, where

$$\vec{V}(\alpha) = \int d\vec{n} \vec{n} p_n(\alpha), \quad (45)$$

and, from (8), the optimal fidelity reads $F = [1 + \sum_{\alpha} V(\alpha)]/2$. Closed expressions of the fidelity for the lowest values of N can be derived from (44) and (45) using

$$\int d\vec{n} n_{i_1} n_{i_2} \cdots n_{i_q} = \frac{1}{K_q} \delta_{\{i_1 i_2 \delta_{i_3 i_4} \cdots \delta_{i_{q-1} i_q\}}, \quad (46)$$

where the normalization factor is $K_q = q!!$ in 2D and $K_q = (q+1)!!$ in 3D, and the indexes in curly brackets are fully symmetrized, e.g., $\delta_{\{i_1 i_2 \delta_{i_3 i_4}\}} = \delta_{i_1 i_2} \delta_{i_3 i_4} + \delta_{i_1 i_3} \delta_{i_2 i_4} + \delta_{i_1 i_4} \delta_{i_2 i_3}$. Obviously, the integral (46) vanishes for q odd. For larger values of N the expressions became rather involved and we have resorted to a numerical calculation.

The 2D case is illustrated in Fig. 1, where the average fidelity for the above two guesses and N in the range 10–60 is shown. In both instances the fidelity approaches unity as N increases, and the OG always performs better than the CLG, as it should. Notice that to make the graphs more easily readable we have interpolated between integer points.

At this point, it is convenient to define the scaled error function

$$\epsilon_N = N(1 - F), \quad (47)$$

and the limit

$$\epsilon = \lim_{N \rightarrow \infty} \epsilon_N, \quad (48)$$

which gives the first order coefficient of the fidelity in the large N expansion, $F = 1 - \epsilon/N + \cdots$ (the asymptotic behavior will be properly discussed in Sec. IV). Fig. 2 shows ϵ_N as a function of N for 2D states. One read-




FIG. 2: Scaled error $\epsilon_N = N(1 - F)$ in the 2D case for collective (solid line), OG (dot-dashed line), CLG (dashed line) and greedy (diamonds) schemes.




FIG. 3: Scaled error $\epsilon_N = N(1 - F)$ in the 3D case for collective (solid line), OG (dot-dashed line), CLG (dashed line) and greedy (diamonds) schemes.

ily sees that the CLG gives $\epsilon_N^{\text{CLG}} \approx 3/8$ [22], while for collective measurements one has $\epsilon_N^{\text{COL}} \approx 1/4$, in agreement with Eq. (27). The stability of the curves ϵ_N^{COL} and ϵ_N^{CLG} shows that the fidelity is well approximated by $F = 1 - \epsilon/N$ for such small values of N as those in the figure. This asymptotic regime is not yet achieved by the OG, however we will show in Sec. IV that the OG gives $\epsilon_N^{\text{OG}} = 1/4$, thus matching the collective bound for large N .

Fig. 3 shows the scaled error in the 3D case. Again, one readily sees that the OG performs better than the CLG. However, the improvement does not seem to be enough to match the collective bound. We prove analytically in the next section that $\epsilon_N^{\text{OG}} = 13/12 > \epsilon_N^{\text{COL}} = 1$.

In previous paragraphs we have presented the most basic scheme, i.e., that with a minimal number of orientations of the measuring devices and without exploiting classical communication. A next step in complexity is to consider a more general set of fixed directions $\{m_k\}$. It is intuitively clear that, assuming some sort of isotropy, the more directions are taken into account, the better the estimation procedure will be. For instance, in 2D we may consider a set of directions given by the

angles $\theta_k = k\pi/N$, where $k = 1, \dots, N$. The set of outcomes χ can be expressed as an N -digit binary number $\chi = i_N i_{N-1} \dots i_2 i_1$, where $i_k (= 0, 1)$ and the fidelity then reads

$$F = \frac{1}{2} + \frac{1}{2} \sum_{\chi=00\dots 0}^{2^N-1} \left| \int dn \vec{n} \prod_{k=1}^N \frac{1 + (-1)^{i_k} \vec{n} \cdot \vec{m}_k}{2} \right|. \quad (49)$$

Analytical results for low N can be obtained using (46). For large N , numerical computations show that this “isotropic strategy” is indeed better than tomography (see [35] for explicit results), but there is no substantial improvement.

For 3D states, however, it is really important to consider more general fixed measurement schemes, such as a 3D version of the isotropic one we have just discussed. One can readily see from Fig. 3 that the tomographic OG does not saturate asymptotically the collective bound and one could be tempted to think that classical communication may be required to attain it [19, 22]. That would somehow indicate a fundamental difference between the estimation of 2D and 3D states.

There is a difficulty in implementing the isotropic scheme for 3D states since the notion of isotropic distribution of directions is not uniquely defined, which contrasts with the 2D case. A particularly interesting scheme that encapsulates this notion (at least asymptotically) and enables us to perform analytical computations consists of measurements along a set of random directions. With the same notation as in (49), the fidelity for this set of directions can be written as

$$F = \frac{1}{2} + \frac{1}{2} \sum_{\chi=0\dots 0}^{2^N-1} \int \prod_{k=1}^N dm_k \left| \int dn \vec{n} \frac{1 + (-1)^{i_k} \vec{n} \cdot \vec{m}_k}{2} \right|. \quad (50)$$

In Fig. 4 we show the scaled error, ϵ_N , obtained from numerical simulations for rather large N . One readily sees the improvement of the random scheme over the tomographic OG. We will show in the next section that the former indeed attains the collective bound asymptotically, thus resolving the puzzle of whether classical communication is needed or not. A numerical fit gives a value $\epsilon_N^{\text{rand}} = 1.002 \pm 0.008$, which provides a numerical check of the analytical results of section IV below.

B. Adaptive Measurements

In this subsection we will discuss schemes that make use of classical communication. In principle, they should be more efficient than those considered so far.

1. One step adaptive

We first review a method put forward by Gill and Massar [19], which we call “one step adaptive”. This scheme,




FIG. 4: Scaled error ϵ_N of the random scheme (triangles) as compared to the OG (dot-dashed line) and the optimal collective scheme (solid line).

although very simple, suffices to show in a very straightforward way that local measurements attain the collective bounds in 2D and 3D. It also has the nice feature that only one reorientation of the measuring device is required.

The basic idea of the method is to split the measurements in two stages. In the first one a small number of copies is used to obtain a rough estimate $\vec{M}_0$ of the state. In the second stage the remaining copies are used to refine the estimate by measuring on a plane orthogonal to $\vec{M}_0$. This strategy has a clear motivation from the information theory point of view. A measurement can be regarded as a query that one makes to a system. The most informative queries are those for which the prior probabilities of each outcome are the same. Measurements on the orthogonal plane to $\vec{M}_0$ have this feature. The method turns out to be efficient if the number of copies used in each of the two stages is carefully chosen.

To be more concrete, suppose we are given N copies of an unknown qubit state. Let N_0 stand for the number of copies used in the first stage and let $\bar{N} = N - N_0$ stand for the rest. In the 2D (3D) case, one measures $N_0/2$ ($N_0/3$) copies along two (three) fixed orthogonal directions and infers the guess $\vec{M}_0$. In the second stage, one measures $\bar{N}$ ($\bar{N}/2$) along $\vec{u}$ ($\vec{u}, \vec{v}$), which are chosen so that $\{\vec{u}, \vec{v}, \vec{M}_0\}$ is an orthonormal basis, and infers the final guess $\vec{M}$. If $N_0 \propto N^a$ with $0 < a < 1$, one can show that the one step adaptive scheme saturates the collective bound in the asymptotic regime [19]. Although we do not have a rigorous proof, our numerical analysis reveals that the optimal value of a , i.e., the one that gives the maximal fidelity, is $a \sim 1/2$. For other choices of a the scheme can even be less efficient than some fixed measurement schemes. For the benefit of the reader, we present a detailed discussion of the method and a proof of the asymptotic limit within our unified framework in Appendix A.

2. Greedy scheme

We now move forward to more sophisticated schemes and discuss one that exploits much more efficiently classical communication. The idea behind it is to maximize the average fidelity at each single measurement step. It is called “greedy” because it does not take into account the total number of available copies, instead, it treats each copy as if it were the last one [20, 21].

We first need to introduce some notation. Recall that the set of outcomes χ can be expressed as a N -digit binary number $\chi = i_N i_{N-1} \dots i_2 i_1$ ($i_k = 0, 1$). Since we allow the k -th measurement to depend on the list of previous outcomes, $i_{k-1} i_{k-2} \dots i_2 i_1 \equiv \chi_k$ (note that $\chi = \chi_N$), we have $\vec{m}(\chi_k)$ instead of $\vec{m}_k$. This is a compact notation where the length k of the string χ_k denotes the number of copies upon which we have already measured. The orthogonality of the von Neumann measurements is imposed by the constraint

$$\vec{m}(1\chi_{k-1}) = -\vec{m}(0\chi_{k-1}), \quad (51)$$

where $1\chi_{k-1}$ is the list of length k obtained by prepending 1 to the list χ_{k-1} , and similarly for $0\chi_{k-1}$. In general, the number of independent vectors for a given N is $(\sum_{k=1}^N 2^k)/2 = 2^N - 1$. For example, if $N = 2$ there are three independent directions, which can be chosen as $\vec{m}(0), \vec{m}(00), \vec{m}(01)$, and the other three are obtained using Eq. (51). Since the first measurement can be chosen at will, this number is reduced to $2^N - 2$.

The general expression of the conditional probability thus reads

$$p_n(\chi) = \prod_{k=1}^N \frac{1 + \vec{n} \cdot \vec{m}(\chi_k)}{2}, \quad (52)$$

and, as discussed in the introduction, the OG gives a fidelity $F = (1 + \Delta)/2$, where

$$\Delta = \sum_{\chi=00\dots0}^{2^N-1} \left| \int d\vec{n} \vec{n} p_n(\chi) \right|. \quad (53)$$

We could in principle attempt to maximize this expression with respect to *all* the independent variables, i.e., all independent $\{\vec{m}(\chi_k)\}$. However, the maximization process very quickly becomes extremely difficult. In the greedy scheme one takes a more modest approach: one maximizes at each step k . This enables us to find a compact algorithm for computing the fidelity, as we discuss below. Furthermore, we show in Appendix B that in this situation the optimal local measurement at each step is indeed of von Neumann type, i.e., any other POVM will perform worse.

Let us concentrate on the last step, N , of the greedy scheme. Suppose we have optimised the previous $N - 1$ measurements and have obtained a string of outcomes χ_{N-1} . To ease the notation, let us denote the direction of the last measurement by $\vec{m}_N$, namely $\vec{m}_N \equiv$

$\vec{m}(0_{\chi_{N-1}}) = -\vec{m}(1_{\chi_{N-1}})$. We then need to maximize

$$d(\chi_N) = |\vec{V}(0_{\chi_{N-1}})| + |\vec{V}(1_{\chi_{N-1}})|. \quad (54)$$

Here

$$\vec{V}(i_N \chi_{N-1}) = \int dn \vec{n} p_n(\chi_{N-1}) \frac{1 + (-1)^{i_N} \vec{n} \cdot \vec{m}_N}{2}, \quad (55)$$

or, equivalently,

$$\vec{V}(i_N \chi_{N-1}) = \frac{1}{2} [\vec{V}(\chi_{N-1}) + (-1)^{i_N} \mathbf{A}(\chi_{N-1}) \vec{m}_N], \quad (56)$$

where $\mathbf{A}$ is the real positive symmetric matrix with elements

$$A_{kl}(\chi_{N-1}) = \int dn n_k n_l p_n(\chi_{N-1}). \quad (57)$$

Therefore

$$d(\chi_N) = \frac{1}{2} \left\{ |\vec{V}(\chi_{N-1}) + \mathbf{A}(\chi_{N-1}) \vec{m}_N| + |\vec{V}(\chi_{N-1}) - \mathbf{A}(\chi_{N-1}) \vec{m}_N| \right\}. \quad (58)$$

Notice that for 2D states and fixed $d(\chi_N)$ the points $\vec{\mu} = \mathbf{A} \vec{m}_N$ lay on an ellipse with focus at $\pm \vec{V}$ (an ellipsoid for 3D states). In addition they fulfil the normalization constraint

$$\vec{\mu} \cdot (\mathbf{A}^{-2} \vec{\mu}) = 1, \quad (59)$$

which also defines an ellipse (ellipsoid in 3D) centered at the origin. As usual, optimality tells us that the maximum of $d(\chi_N)$ occurs at the points of tangency of the ellipses (ellipsoids). This provides a geometrical procedure for finding the optimal direction $\vec{m}_N$ and an algorithm for computing $|\vec{V}(\chi_N)|$.

We now proceed to obtain some explicit expressions for low N . We discuss only the 3D case, as the 2D case is completely analogous (numerical results for 2D states are shown in Fig. 1 and Fig. 2).

When we only have one copy of the state, $N = 1$, the Bloch vector of the measurement can be chosen in any direction, say $\vec{e}_x$, i.e., $\vec{m}(0) = -\vec{m}(1) = \vec{e}_x$. The explicit computation of the vector $\vec{V}$ in (7) gives

$$\vec{V}(\chi_1) = \frac{1}{6} \vec{m}(\chi_1), \quad (60)$$

and $F = 2/3$, as expected from (37) [or (18)].

The first non-trivial case is $N = 2$. The matrix $\mathbf{A}(\chi_1)$ reads

$$A_{kl}(\chi_1) = \frac{1}{6} \delta_{kl} \quad \chi_1 = 0, 1, \quad (61)$$

i.e., $\mathbf{A}(\chi_1)$ is independent of χ_1 and proportional to the identity. The maximum of (58) occurs for $\vec{m}_2 \perp \vec{e}_x$, so we choose $\vec{m}_2 = \vec{e}_y$, which means, $\vec{m}(00) = \vec{m}(01) = \vec{e}_y$, (notice that in general these two vectors do not need to be

equal, they are only required to be orthogonal to $\vec{m}(0)$). Because of (51), we also have $\vec{m}(10) = \vec{m}(11) = -\vec{e}_y$. The OG reads

$$\vec{M}^{(2)}(\chi) = \frac{\vec{m}(\chi_2) + \vec{m}(\chi_1)}{\sqrt{2}}, \quad (62)$$

e.g., $\vec{M}^{(2)}(01) = [\vec{m}(01) + \vec{m}(1)]/\sqrt{2} = [\vec{m}(01) - \vec{m}(0)]/\sqrt{2} = [\vec{e}_y - \vec{e}_x]/\sqrt{2}$. One obtains

$$|\vec{V}(\chi_2)| = \frac{\sqrt{2}}{12} \quad (63)$$

for all χ_2 , which implies

$$F^{(2)} = \frac{3 + \sqrt{2}}{6}. \quad (64)$$

The case $N = 3$ can be computed along the same lines. One can easily see that $\vec{m}(\chi_3)$ has to be perpendicular to $\vec{m}(\chi_2)$ and $\vec{m}(\chi_1)$. This shows that, up to $N = 3$, the greedy approach does not use classical communication, i.e., the directions of the measuring devices are only required to be mutually orthogonal, independently of the outcomes. The optimal guess is a straightforward generalization of (62):

$$\vec{M}^{(3)}(\chi) = \frac{\vec{m}(\chi_3) + \vec{m}(\chi_2) + \vec{m}(\chi_1)}{\sqrt{3}}, \quad (65)$$

and the fidelity reads

$$F^{(3)} = \frac{3 + \sqrt{3}}{6}. \quad (66)$$

The above results could have been anticipated. As already mentioned, the outcomes of a measurement on the plane orthogonal to the guess have roughly the same probability and are, hence, most informative. One can regard these measurements as corresponding to mutually unbiased observables, i.e., those for which the overlap between states of different basis (related to each observable) is constant [36]. Hence, there is no redundancy in the information about the state acquired from the different observables. This point of view also allows to extend the notion of (Bloch vector) orthogonality to states in spaces of arbitrary dimension.

The case $N = 4$ is even more interesting, since four mutually orthogonal vectors cannot fit onto the Bloch sphere. We expect classical communication to start playing a role here. Indeed the Bloch vectors $\vec{m}(\chi_4)$ do depend on the outcomes of previous measurements. They can be compactly written as [40]

$$\vec{m}(\chi_4) = \frac{(-1)^{i_4}}{\sqrt{2}} \sum_{k=1}^2 \vec{m}(\chi_k) \times \vec{m}(\chi_3). \quad (67)$$

Again, one can see that the vectors $\vec{m}(\chi_4)$ are orthogonal to the guess one would have made with the first three measurements. The fidelity in this case is

$$F^{(4)} = \frac{15 + \sqrt{91}}{30}. \quad (68)$$

For larger N , we have computed the fidelity of the greedy scheme by numerical simulations. In Fig 3 (Fig. 1 and Fig. 2 for 2D states) we show the results for $10 \leq N \leq 60$ (diamonds). Notice that the greedy scheme is indeed better than fixed measurement schemes and approaches the collective bound very fast.

Actually, the greedy scheme is the best we can use if we do not know a priori the number of copies that will be available; obviously, the best one can do in these circumstances is to optimise at each step. However, if N is known, we have extra information that some efficient schemes could exploit to increase the fidelity. We next show that this is indeed the case.

3. General LOCC scheme

In the most general LOCC scheme one is allowed to optimise over all the Bloch vectors $\{m(\chi_k)\}$, thus taking into account the whole history of outcomes. Up to $N = 3$ the results are the same as for the greedy scheme: orthogonal Bloch vectors for the measurements and no classical communication required. The results (64) and (66) are, therefore, the largest fidelity that can be attained by any LOCC scheme.

The most interesting features appear at $N = 4$. Here there are 14 independent vectors, which can be grouped into two independent families of seven vectors. With such a large number of vectors an analytical calculation is too involved and we have resorted partially to a numerical optimization. The solution exhibits some interesting properties. First, one obtains that $\vec{m}(\chi_1) \perp \vec{m}(\chi_2)$, for all χ_1 and χ_2 , as in the $N = 2$ and $N = 3$ cases. Therefore one can choose $\vec{m}(\chi_1) = (-1)^{i_1} \vec{e}_x$ and $\vec{m}(\chi_2) = (-1)^{i_2} \vec{e}_y$. Only for the third and fourth measurement one really has to take different choices in accordance to the sequence of the preceding outcomes. The Bloch vectors of the third measurement can be parametrized by a single angle α as

$$(-1)^{i_3} \vec{m}(\chi_3) = \cos \alpha \vec{u}_1(\chi_2) + \sin \alpha \vec{v}_1(\chi_2), \quad (69)$$

where

$$\begin{aligned} \vec{u}_1(\chi_2) &= \vec{m}(\chi_1) \times \vec{m}(\chi_2), \\ \vec{v}_1(\chi_2) &= \vec{u}_1(\chi_2) \times \vec{s}(\chi_2), \\ \vec{s}(\chi_2) &= \frac{\vec{m}(\chi_2) + \vec{m}(\chi_1)}{\sqrt{2}}. \end{aligned} \quad (70)$$

Notice that, rather unexpectedly, $\vec{m}(\chi_1)$, $\vec{m}(\chi_2)$ and $\vec{m}(\chi_3)$ are not mutually orthogonal. The optimal value of this angle is $\alpha = 0.502$. Although we cannot give any insight as to why this value is optimal, in agreement with our intuition one sees that $\vec{m}(\chi_3) \perp \vec{M}(\chi_2)$, i.e., the third measurement probes the plane orthogonal to the Bloch vector one would guess from the first two outcomes [see Eq. (62)]. The vectors of the fourth measurement can be parametrized by two angles, β and γ , as

$$(-1)^{i_4} \vec{m}(\chi_4) = \cos \gamma \vec{u}_2(\chi_3) + \sin \gamma \vec{v}_2(\chi_3), \quad (71)$$

where

$$\begin{aligned} \vec{u}_2(\chi_3) &= \vec{s}(\chi_2) \times \vec{m}(\chi_3), \\ \vec{v}_2(\chi_3) &= \cos \beta \vec{m}(\chi_3) - \sin \beta \vec{s}(\chi_2) \end{aligned} \quad (72)$$

The optimal values of these angles are $\beta = 0.584$, $\gamma = 0.538$, and the corresponding fidelity is $F_{\text{general}}^{(4)} = 0.8206$. This is just 1.5% lower than the absolute bound $5/6 = 0.8333$ attained with collective measurement, Eq. (37). Note that this value is slightly larger than the fidelity obtained with the greedy scheme, $F_{\text{general}}^{(4)} > F_{\text{greedy}}^{(4)} = (15 + \sqrt{91})/30 \approx 0.8180$. The extra information consisting of the number of available copies has indeed been used to attain a larger fidelity. We conclude that for $N > 3$, it pays to relax optimality at each step, and greedy schemes [20, 21] are thus not optimal. We would like to remark that if, for some reason, some copies are lost or cannot be measured, the most general scheme will not be optimal, since it has been designed for a specific number of copies. We have also computed the values of the maximal LOCC fidelities for $N = 5, 6$: $F_{\text{general}}^{(5)} = 0.8450$ and $F_{\text{general}}^{(6)} = 0.8637$. Beyond $N = 6$ the small differences between this and the greedy scheme become negligible.

IV. LOCAL SCHEMES IN THE ASYMPTOTIC LIMIT

The asymptotic expression of the fidelity enables us to compare different schemes independently of the number of copies. If two schemes have the same asymptotic fidelity, it is justified to say that they have the same efficiency, and conversely. Here we will compute such asymptotic expansions. We will show that, asymptotically, classical communication is not needed to attain the absolute upper bound given by the maximum fidelity of the most general collective measurements. Some of the results presented in this section were obtained by two of the authors by explicit computations in [22]. Here we will use a statistical approach that relate the Fisher information I [37] with the average fidelity F . This approach will greatly simplify our earlier derivations.

We label the independent state parameters by the symbol η . This symbol will refer to the two angles θ, ϕ for 3D states: $\eta \equiv (\theta, \phi)$; and the polar angle θ for 2D states: $\eta \equiv \theta$.

Assume that under a sensible measurement and estimation scheme (we mean by that a scheme that leads to a perfect determination of the state when $N \rightarrow \infty$, i.e., $F \xrightarrow{N \rightarrow \infty} 1$) the estimated state is close to the signal state, that is, their respective parameters $\hat{\eta}(\chi)$ and η differ by a small amount. In this Section, a hat ($\hat{\cdot}$) will always refer to estimated parameters, the fidelity $f_n(\chi)$, Eq. (2), will be denoted by $f_n(\hat{\eta})$, and similarly, the probability $p_n(\chi)$ will be written as $p_n(\hat{\eta})$. Note that the guessed parameters $\hat{\eta}(\chi)$ are based on a particular outcome χ .

This dependence will be implicitly understood when no confusion arises.

The fidelity can be approximated by the first terms of its series expansion:

$$f_\eta(\hat{\eta}) \approx 1 + \frac{1}{2} \frac{\partial^2 f}{\partial \hat{\eta}_i \partial \hat{\eta}_j} \Big|_{\hat{\eta}=\eta} (\hat{\eta}_i - \eta_i)(\hat{\eta}_j - \eta_j), \quad (73)$$

where we have used that $f_\eta(\eta) = 1$ and $\partial f_\eta / \partial \hat{\eta}|_{\hat{\eta}=\eta} = 0$. Averaging over all possible outcomes, we have

$$F(\eta) \approx 1 + \frac{1}{2} \text{tr} [\mathbf{H}(\eta) \mathbf{V}(\eta)], \quad (74)$$

where

$$F(\eta) \equiv \sum_{\chi} p_\eta(\chi) f_\eta[\hat{\eta}(\chi)], \quad (75)$$

is the “pointwise” fidelity, $\mathbf{H}(\eta)$ is the Hessian matrix of $f_\eta(\hat{\eta})$ at $\hat{\eta} = \eta$, and $\mathbf{V}(\eta)$ is the covariance matrix, with elements $V_{ij}(\eta) = \sum_{\chi} p_\eta(\chi) (\hat{\eta}_i - \eta_i)(\hat{\eta}_j - \eta_j)$.

It is well known that the variance of an unbiased estimator is bounded by

$$\mathbf{V}(\eta) \geq \frac{1}{\mathbf{I}(\eta)}, \quad (76)$$

the so called Cramér-Rao bound [19, 25, 38], where the Fisher information matrix $\mathbf{I}(\eta)$ is defined as

$$I_{ij}(\eta) = \sum_{\chi} p_\eta(\chi) \frac{\partial \ln p_\eta(\chi)}{\partial \eta_i} \frac{\partial \ln p_\eta(\chi)}{\partial \eta_j}. \quad (77)$$

The conditional probability $p_\eta(\chi)$ regarded as a function of η is called the likelihood function $\mathcal{L}(\eta) = p_\eta(\chi)$. It is also well known that the bound (76) is attained by the maximum likelihood estimator (MLE) [39], defined as $\hat{\eta}^{\text{MLE}} = \text{argmax } \mathcal{L}(\eta)$. Hence this bound is tight.

A link between the Fisher information and the fidelity is obtained by combining (74) and (76), and noticing that $\mathbf{H}(\eta)$ is negative definite. We thus have

$$F(\eta) \leq 1 + \frac{1}{2} \text{tr} \frac{\mathbf{H}(\eta)}{\mathbf{I}(\eta)} \quad (78)$$

to leading order and for any unbiased estimation scheme.

The Fisher information is additive. This means that if $p_\eta^{(2)}(\chi, \chi') = p_\eta(\chi) p'_\eta(\chi')$, which happens when we perform two measurements [say, $\{O(\chi)\}$ and $\{O'(\chi')\}$] on two identical states, the Fisher information of the combined measurement is simply $\mathbf{I}^{(2)}(\eta) = \mathbf{I}(\eta) + \mathbf{I}'(\eta)$. In particular, for N identical measurements, we have $\mathbf{I}^{(N)}(\eta) = N \mathbf{I}(\eta)$.

Finally, since the OG is a better estimator, and it is asymptotically unbiased, we must have

$$F^{\text{OG}}(\eta) \simeq F^{\text{MLE}}(\eta) = 1 + \frac{1}{2N} \text{tr} \frac{\mathbf{H}(\eta)}{\mathbf{I}(\eta)} \quad (79)$$

to leading order, where the fidelities refer to an estimation scheme consisting of N identical measurements. Below we use Eq. (79) to compute the asymptotic limits of the fixed measurement schemes discussed in Sec. III A.

We would like to remark that the Cramér-Rao bound assumes some regularity conditions on the figure of merit and the estimators. These conditions are satisfied for the problem considered here, but the bound may not hold in more general situations such as the estimation of a mixed qubit state (see [23]).

A. 2D states

The 2D case is rather simple because the states have just one parameter and the Fisher information is a single number. Moreover, *any* von Neumann measurement whose vector lies on the equator of the Bloch sphere performed on a 2D system has $\mathbf{I} = 1$, as can be checked by plugging

$$p_\theta(\pm 1) = \frac{1 \pm \cos(\theta - \theta_m)}{2} \quad (80)$$

into (77), where θ_m is the polar angle of $\vec{m}$, the direction along which the von Neumann measurement is performed. Therefore, in 2D the Fisher information for a set of N measurements (identical or not) is $\mathbf{I}^{(N)} = N$.

Since the Hessian is

$$\mathbf{H} = \frac{\partial^2 f}{\partial \theta^2} \Big|_{\hat{\theta}=\theta} = -\frac{1}{2}, \quad (81)$$

any sensible local measurement scheme on 2D states will yield

$$F(\theta) = 1 - \frac{1}{4N} + \dots \quad (82)$$

Note that this fidelity is independent of θ , so it coincides with the average fidelity $F = \int d\theta F(\theta)/(2\pi) = 1 - 4/N$.

We recall that this fidelity is attained by the MLE, and hence also by the OG. We note that it coincides with the collective bound, Eq. (27). This implies that the collective bound is attained by tomography, without classical communication. It is quite surprising that the most basic scheme, with measurements along two fixed orthogonal directions, saturates already the collective bound asymptotically.

B. 3D states

The 3D case is more involved. The results shown in Fig. 3 hint that tomography does not saturate the collective bound. There are, however, other measurement schemes that do saturate this bound and still do not require classical communication. We prove these two statements below.

1. Tomography

Tomography, as explained in Sec. III A, consists in measuring along three orthogonal directions on the Bloch sphere. We will concentrate on the OG estimator, or more precisely on the MLE, which is asymptotically equivalent. We do not discuss the CLG, Eq. (43), as it does not attain the collective bound even for 2D states (see Fig. 2).

We first compute the Fisher information. Consider a scheme that consists in repeating N times the following: take 3 copies of the state and perform a measurement along $\vec{e}_x$ on the first copy, along $\vec{e}_y$ on the second copy, and along $\vec{e}_z$ on the third copy (recall that $N = 3N$). These three von Neumann measurements can be regarded as a single measurement with 2^3 possible outcomes labeled by $\chi = (\chi_1, \chi_2, \chi_3)$, where $\chi_j = \pm 1$. The probability of obtaining an outcome χ is

$$p_\eta(\chi) = \prod_{j=x,y,z} \frac{1}{2} (1 + \chi_j \vec{n} \cdot \vec{e}_j). \quad (83)$$

The Fisher information matrix $I(\theta, \phi)$ of such elementary measurement is obtained by substituting Eq. (83) in Eq. (77). Note that the Fisher information of this scheme is $I^{(N)}(\theta, \phi) = NI(\theta, \phi)$.

The Hessian of the fidelity is

$$H(\theta, \phi) = - \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1 - \cos 2\theta}{4} \end{pmatrix}, \quad (84)$$

which turns out to be independent of ϕ . With this, we obtain

$$\frac{1}{N} \text{tr} \frac{H(\theta, \phi)}{I(\theta, \phi)} = \frac{3}{16N} \times \frac{35 + 28 \cos 2\theta + \cos 4\theta - 8 \cos 4\phi \sin^4 \phi}{9 + 7 \cos 2\theta - 2 \cos 4\phi \sin^2 \theta}. \quad (85)$$

Integrating over the isotropic prior probability dn , Eq. (31), we obtain

$$\frac{1}{N} \int dn \text{tr} \frac{H(\theta, \phi)}{I(\theta, \phi)} = -\frac{13}{18N}. \quad (86)$$

Recalling (79) and $N = 3N$ we finally get

$$F^{\text{OG}} = 1 - \frac{13}{12N} + \dots \quad (87)$$

As mentioned above, the collective bound $F = 1 - 1/N$ is not attained by tomography. At this point, the question arises whether classical communication is necessary to attain this bound. We next show that this is not the case.

2. Random scheme

We now consider the so called random scheme i.e, a scheme in which measurements are performed along random directions chosen from an isotropic distribution. In contrast to tomography, which only takes into account three fixed directions, this scheme explores the Bloch sphere isotropically if a large number of copies is available. Therefore one can expect that it will perform much better.

This approach is equivalent to performing a covariant (continuous) POVM on each one of the copies separately. Here, we instead regard it as von Neumann measurements and a classical ancilla, e.g., a “roulette”, that tells us along which direction we measure. From this point of view the outcome parameters are given by $\chi = (\xi, (u \equiv \cos \vartheta), \varphi)$, where ϑ and φ are the azimuthal and polar angles of the direction $\vec{m}(u, \varphi)$ of the measurement, and $\xi = \pm 1$ is the corresponding outcome.

Let us compute the Fisher information for a specific state $\eta = ((v \equiv \cos \theta), \phi)$. Since this strategy is isotropic, the pointwise fidelity $F(\eta)$ is independent of η , and we conveniently choose $\eta = ((v = 0), 0) = \underline{0}$. By the same argument, no average over η will be needed: $F = F(\eta)$. The probability is given by

$$p_\eta(\chi) = \frac{1 + \xi \vec{n} \cdot \vec{m}(u, \varphi)}{2}, \quad (88)$$

and the Fisher information reads

$$I_{ij}(\eta) = \sum_{\xi=\pm 1} \int \frac{du d\varphi}{4\pi} p_\eta(\chi) \frac{\partial \ln p_\eta(\chi)}{\partial \eta_i} \frac{\partial \ln p_\eta(\chi)}{\partial \eta_j}. \quad (89)$$

The diagonal elements read

$$I_{vv}(\underline{0}) = \frac{1}{8\pi} \sum_{\xi=\pm 1} \int \frac{u^2 du d\varphi}{1 + \xi \sqrt{1 - u^2} \cos \varphi} = \frac{1}{2}, \quad (90)$$

$$I_{\phi\phi}(\underline{0}) = \frac{1}{8\pi} \sum_{\xi=\pm 1} \int \frac{(1 - u^2) \sin^2 \varphi du d\varphi}{1 + \xi \sqrt{1 - u^2} \cos \varphi} = \frac{1}{2}. \quad (91)$$

As for the off-diagonal elements, a straightforward calculation gives

$$I_{v\phi}(\underline{0}) = I_{\phi v}(\underline{0}) = 0, \quad (92)$$

as one could expect, since gaining information on v does not provide information on ϕ and viceversa. The Fisher information matrix thus reads

$$I(\underline{0}) = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (93)$$

The Hessian of the fidelity is

$$H_{ij}(\underline{0}) = \left. \frac{\partial^2 f}{\partial \hat{\eta}_i \partial \hat{\eta}_j} \right|_{\hat{\eta}=\underline{0}} = -\frac{\delta_{ij}}{2}. \quad (94)$$

Finally, using (79) we obtain

$$F^{\text{OG}} = 1 - \frac{1}{N} + \dots \quad (95)$$

We conclude that asymptotically classical communication is not required to saturate the collective bound: a measurement scheme based on a set of random directions does the job.

V. CONCLUSIONS

We have presented a selfcontained and detailed study of several estimation schemes when a number N of identical copies of a qubit state is available. We have used the fidelity as a figure of merit and presented a general framework which enables us to treat collective as well as local measurements on the same footing.

We have considered two interesting situations: that of a completely unknown qubit state (3D case), and that of a qubit laying on the equator of the Bloch sphere (2D case). We have obtained the optimal measurements and maximum fidelities for the most general collective strategies. These results, although well known, were scattered in the literature and rederived several times. Here we have obtained them within a direct and unified framework. The solution in the 2D case is strikingly simple, and can be extended to the case of optimal covariant phase estimation.

These collective schemes yield the ultimate fidelity bounds that can be achieved by any scheme, thus setting a natural scale of what is a good or a bad estimation. However, they require a joint measurement on the N copies, which is usually very difficult, if not impossible, to be implemented in practice. The main part of this paper has therefore focused on measurements that can be implemented in a laboratory with nowadays technology: local von Neumann measurements.

In the 2D case we have shown that, quite surprisingly, the most basic tomographic scheme, i.e., measurements along two fixed orthogonal directions with the adequate data processing (the OG), gives already a fidelity that is asymptotically equal to the collective bound. We have obtained this limit using the Cramér-Rao bound, which is particularly simple to compute in this situation. This result is in agreement with the direct (and lengthier) computation presented in [22].

For the 3D states, tomography, i.e., measurements along three fixed orthogonal directions, fails to give the asymptotic collective bound, even with the best data processing. The main reason of this failure is that the Bloch sphere is not explored thoroughly. We have considered an extension that is asymptotically isotropic: a series von Neumann measurements along random directions. We have proved that this scheme, which does not make use of classical communication, does saturates the collective bound. Hence, we conclude that in the large N limit, an

estimation procedure based on local measurements without classical communication does perform as well as the most efficient and sophisticated collective schemes.

We have also discussed local schemes with classical communication, i.e., schemes in which the measurements are devised in such a way that they take into account previous outcomes. We have studied in detail the one step adaptive scheme of Gill and Massar [19], which has very interesting features: only two measurement orientations are required, adaptivity is only used once, and the estimation is made by a CLG, which can be read off from the frequency of outcomes. The economy of resources in this scheme may raise doubts about its efficiency. In Appendix A we give a simple proof that for large N it indeed attains the collective bounds.

We have also studied strategies that make a more intensive use of classical communication. In the greedy scheme optimization is performed at each measurement step [20, 21]. This scheme is the best approach one can take if the actual number of available copies is not known. We have given a geometrical condition for sequentially finding the optimal measurements and have proved that they have to be of von Neumann type (see App. B), i.e., no general local POVM's will perform better in this context. We have illustrated the performance of the method with numerical simulations and have shown the behaviour of the optimal collective scheme is reached for very low values of N . This occurs for N as low as $N = 20$ in 2D and slightly above, $N = 45$, in 3D.

In the most general scheme we see that up to $N = 3$ ($N = 2$ in 2D) there is no need for classical communication: the optimal measurements correspond to a set of mutually unbiased observables. For larger N , the knowledge of the actual value of N provides an extra information that translates into an increase of the fidelity. From the practical point of view, however, this difference is negligible already at the level of a few copies ($N \gtrsim 6$).

Our approach may be extended to other situations. For instance, the problem of estimating (qubit) mixed states, which is much more involved, can be tackled along the lines described here [23]. It would also be interesting to consider qudits and check whether a set of mutually unbiased observables provides the optimal local estimation scheme when the number of copies coincides with the number of independent variables that parametrize the qudit state.

Acknowledgments

It is a pleasure to thank M. Baig for his collaboration at early stages of this work. We also thank R. Gill and M. Ballester for useful discussions. We acknowledge financial support from Spanish Ministry of Science and Technology project BFM2002-02588, CIRIT project SGR-00185, and QUPRODIS working group EEC contract IST-2001-38877.

APPENDIX A

The simplest LOCC approach is exemplified by the “one step adaptive” scheme [19]. Measurements are performed along just two different directions, and CLG is used [41]. The scheme has two stages and classical communication is used only once, in going from the first stage to the second. This is therefore, a very economical scheme from the practical and theoretical point of view. We here review the method and give a straightforward and comprehensive proof that it saturates the collective bound for large N . Given the economy of the scheme, this is not an obvious result at all. We focus only on the 3D case, as the simpler 2D case can be worked out along the same lines.

First stage: One performs $N_0 = N^a$ ($0 < a < 1$) measurements with a sensible estimator, in the sense of Sec. IV, and obtains an estimation $\vec{M}_0$ with a fidelity F_0 :

$$F_0 = \sum_{\chi_0} \int dn \frac{1 + \vec{n} \cdot \vec{M}_0(\chi_0)}{2} p_n(\chi_0), \quad (\text{A1})$$

where χ_0 stands for the list of outcomes obtained in this first stage.

Second Stage: At this point we use the CLG on the remaining $2N \equiv \bar{N} = N - N_0$ copies by measuring along *two* perpendicular directions, $\vec{u}$ and $\vec{v}$, on the plane orthogonal to $\vec{M}_0$. In this basis the final guess can be written as

$$\vec{M} = \vec{M}_0(\chi_0) \cos \omega + (\vec{u} \cos \tau + \vec{v} \sin \tau) \sin \omega. \quad (\text{A2})$$

This parametrization ensures that $\vec{M}$ is unitary. The angles ω and τ depend on the outcomes of this second stage, which are the frequencies $\alpha_i N$, $(1 - \alpha_i)N$, ($i = u, v$). The probabilities are given by $p_n(\alpha)$ in (42), with $n_u = \vec{n} \cdot \vec{u}$ and $n_v = \vec{n} \cdot \vec{v}$. As argued above, we measure on the plane orthogonal to $\vec{M}_0$ because the two outcomes of each measurement have roughly the same probability, $\alpha_i \approx 1/2$, and they are most informative. It is convenient to define the two dimensional vector $\vec{r}$, with components:

$$r_i \equiv 2\alpha_i - 1, \quad i = u, v, \quad (\text{A3})$$

which, on average, is close to $\vec{0}$. This vector gives an estimation of the projection of the signal Bloch vector $\vec{n}$ on the measurement plane (uv plane). Hence, ω is expected to be small ($\vec{M}_0 \approx \vec{M}$) and we make the ansatz

$$\omega = \lambda \sqrt{r_u^2 + r_v^2}, \quad \tan \tau = \frac{r_v}{r_u}, \quad (\text{A4})$$

where the positive parameter λ will be determined later.

The final fidelity for a signal state $\vec{n}$ and outcomes $(\chi_0, \vec{r})$ is

$$f_n(\chi_0, \vec{r}) = \frac{1 + \vec{n} \cdot \vec{M}(\chi_0, \vec{r})}{2}, \quad (\text{A5})$$

and the average fidelity F reads

$$F = \sum_{\chi_0, \vec{r}} \int dn \frac{1 + \vec{n} \cdot \vec{M}(\chi_0, \vec{r})}{2} p_n(\chi_0) p_n(\vec{r}|\chi_0). \quad (\text{A6})$$

Notice that the probability of obtaining the outcome $\vec{r}$, namely, $p_n(\vec{r}|\chi_0)$ [$\equiv p_n(\alpha)$ in (42) with $i = u, v$], is conditioned on χ_0 through the dependence of the second stage measurements on $\vec{M}_0(\chi_0)$.

Since we will compute different averages over χ_0 , $\vec{r} = (r_u, r_v)$ and $\vec{n}$, it is convenient to introduce the following notation:

$$\langle f \rangle_0 = \sum_{\chi_0} f_n(\chi_0, \vec{r}) p_n(\chi_0), \quad (\text{A7})$$

$$\langle f \rangle_r = \sum_{\vec{r}} f_n(\chi_0, \vec{r}) p_n(\vec{r}|\chi_0), \quad (\text{A8})$$

$$\langle f \rangle_n = \int dn f_n(\chi_0, \vec{r}), \quad (\text{A9})$$

and similarly for averages of other functions of χ_0 , $\vec{r}$ and $\vec{n}$. We will denote composite averaging by simply combining subscripts (i.e. $\langle \langle F \rangle_r \rangle_0 \equiv \langle F \rangle_{r,0}$). Therefore, we write

$$F \equiv \langle f \rangle_{r,0,n} \equiv \langle f \rangle. \quad (\text{A10})$$

Since $F = (1 + \Delta)/2$, we have

$$\Delta = \langle \vec{n} \cdot \vec{M} \rangle. \quad (\text{A11})$$

In the expansions that we perform below, we keep only the terms that contribute to the fidelity up to order $1/N$. Recalling that ω is expected to be small, it follows that

$$\cos \omega = 1 - \frac{\lambda^2}{2} [r_u^2 + r_v^2], \quad (\text{A12})$$

$$\sin \omega \cos \tau = \lambda r_u, \quad (\text{A13})$$

$$\sin \omega \sin \tau = \lambda r_v, \quad (\text{A14})$$

to leading order. Therefore, the expectation value in (A11) can be written as

$$\Delta = \left\langle \left(1 - \frac{\lambda^2}{2} \langle r_u^2 + r_v^2 \rangle_r \right) \vec{n} \cdot \vec{M}_0 \right\rangle_{0,n} \quad (\text{A15})$$

$$+ \lambda \left\langle \langle r_u \rangle_r n_u + \langle r_v \rangle_r n_v \right\rangle_{0,n}. \quad (\text{A16})$$

Since r_u, r_v (or equivalently α_u, α_v) are binomially distributed, one readily sees that

$$\langle r_i \rangle_r = n_i, \quad (\text{A17})$$

$$\langle r_i^2 \rangle_r = n_i^2 + \frac{1 - n_i^2}{N}. \quad (\text{A18})$$

We further recall that $\vec{n}$ is unitary and that $\{\vec{M}_0, \vec{u}, \vec{v}\}$ is an orthonormal basis, hence $n_u^2 + n_v^2 = 1 - (\vec{n} \cdot \vec{M}_0)^2$,

and (A16) can be cast as

$$\Delta = \lambda + \left[1 - \frac{\lambda^2}{2} \left(1 + \frac{1}{N} \right) \right] \langle \vec{n} \cdot \vec{M}_0 \rangle_{0,n} - \lambda \langle (\vec{n} \cdot \vec{M}_0)^2 \rangle_{0,n} + \frac{\lambda^2}{2} \left(1 - \frac{1}{N} \right) \langle (\vec{n} \cdot \vec{M}_0)^3 \rangle_{0,n}. \quad (\text{A19})$$

To compute the moments $\langle (\vec{n} \cdot \vec{M}_0)^q \rangle_{0,n}$, we consider the angle δ between $\vec{n}$ and $\vec{M}_0$, which is also expected to be small. We have

$$2F_0 - 1 = \langle \vec{n} \cdot \vec{M}_0 \rangle_{0,n} = \langle \cos \delta \rangle_{0,n} \simeq 1 - \frac{\langle \delta^2 \rangle_{0,n}}{2}, \quad (\text{A20})$$

where we have used (A1). Therefore

$$\begin{aligned} \langle (\vec{n} \cdot \vec{M}_0)^q \rangle_{0,n} &= \langle \cos^q \delta \rangle_{0,n} \simeq 1 - \frac{q}{2} \langle \delta^2 \rangle_{0,n} \\ &= 1 - 2q(1 - F_0). \end{aligned} \quad (\text{A21})$$

Now we plug this result back into (A19) to obtain

$$F = \langle f \rangle = 1 - (1 - \lambda)^2 (1 - F_0) - \frac{1 - 4(1 - F_0)}{2N} \lambda^2. \quad (\text{A22})$$

Since the term $(1 - \lambda)^2 (1 - F_0)$ is always positive, the maximum fidelity is obtained with the choice $\lambda = 1$, and we are left with

$$F = 1 - \frac{1 - 4(1 - F_0)}{2N}. \quad (\text{A23})$$

Since the first estimation is asymptotically unbiased, $1 - F_0$ vanishes for large N_0 (i.e., for large N) and

$$F \simeq 1 - \frac{1}{2N}. \quad (\text{A24})$$

Recalling that $N = (N - N^a)/2$, we finally have

$$F = 1 - \frac{1}{N} + \dots \quad (\text{A25})$$

This concludes the proof.

APPENDIX B

In this appendix we prove that in the greedy scheme the optimal individual measurements on each copy are of von Neumann type. We sketch the proof for 2D states. The 3D case can be worked out along the same lines.

The history of outcomes will be denoted, as usual, by χ . Notice that here we consider general local measurements (local POVMs) with R outcomes, where R is possibly larger than two. Therefore χ is a N -digit integer number in base R : $\chi = i_N i_{N-1} \dots i_1$ ($i_k = 0, 1, \dots, R-1$). As in Sec. IIIB we use the notation $\chi_k = i_k i_{k-1} \dots i_1$. A measurement on the k -th copy

is defined by a set of non-negative rank-one operators $\{O(\chi_k)\}_{i_k=0}^{R-1} = \{O(i_k \chi_{k-1}) \mid i_k = 0, 1, \dots, R-1\}$, where

$$O(\chi_k) = c(\chi_k) [1 + \vec{m}(\chi_k) \cdot \vec{\sigma}]. \quad (\text{B1})$$

The non-negative constants $c(\chi_k)$ and the vectors $\vec{m}(\chi_k)$ are subject to the constraints

$$\sum_{i_k=0}^{R-1} c(\chi_k) = 1, \quad (\text{B2})$$

$$\sum_{i_k=0}^{R-1} c(\chi_k) \vec{m}(\chi_k) = 0, \quad (\text{B3})$$

$$|\vec{m}(\chi_k)| = 1, \quad (\text{B4})$$

which ensures that $O(\chi_k) \geq 0$ and $\sum_{i_k} O(\chi_k) = \mathbb{1}$. Note that we allow $c(\chi_k)$ to be zero, thus taking into account the possibility that each local POVM may have a different number of outcomes without letting R depend on k .

Assume we have measured all but the last copy and we wish to optimise the last measurement. Recall from Sec. I that the fidelity can be written as $F = (1 + \Delta)/2$, where

$$\Delta = \sum_{\chi} |\vec{V}(\chi)|, \quad \vec{V}(\chi) = \int dn \vec{n} p_n(\chi). \quad (\text{B5})$$

To simplify the notation, let us define $r \equiv i_N$, $\vec{m}_r = \vec{m}(r \chi_{N-1})$, and $c_r = c(r \chi_{N-1})$. Then,

$$p_n(\chi) = p_n(\chi_{N-1}) [c_r (1 + \vec{n} \cdot \vec{m}_r)] \quad (\text{B6})$$

and

$$\Delta = \sum_{\chi_{N-1}} \sum_r |\vec{V}(r \chi_{N-1})| = \sum_{\chi_{N-1}} d(\chi_{N-1}), \quad (\text{B7})$$

where we have defined $d(\chi_{N-1})$ as

$$d(\chi_{N-1}) \equiv \sum_r c_r \left| \int dn \vec{n} p_n(\chi_{N-1}) (1 + \vec{n} \cdot \vec{m}_r) \right|. \quad (\text{B8})$$

We further write $\vec{V} \equiv \vec{V}(\chi_{N-1})$ and define the symmetric positive matrix

$$A_{ij} \equiv A_{ij}(\chi_{N-1}) \equiv \int dn n_i n_j p_n(\chi_{N-1}). \quad (\text{B9})$$

Eq. (B8) becomes

$$d = \sum_r c_r |\vec{V} + A \vec{m}_r| \quad (\text{B10})$$

(Hereafter the dependency on χ_{N-1} will be implicitly understood to simplify the notation).

Our task is to maximize (B10). Introducing the Lagrange multipliers λ , $\vec{\gamma}$, and ω_r , the function we need to maximize is actually

$$L = d - \lambda \Lambda - \vec{\gamma} \cdot \vec{\Gamma} - \sum_r \omega_r \Omega_r, \quad (\text{B11})$$

where the constraints

$$\Lambda = \sum_r c_r - 1, \quad (\text{B12})$$

$$\vec{\Gamma} = \sum_r c_r \vec{m}_r, \quad (\text{B13})$$

$$\Omega_r = \frac{\vec{m}_r^2 - 1}{2}, \quad (\text{B14})$$

can be read off from (B2), (B3) and (B4). The factor two in the last expression is introduced for later convenience. Variations with respect to c_r yield

$$\frac{\delta L}{\delta c_r} = |\vec{V} + A\vec{m}_r| - \vec{\gamma} \cdot \vec{m}_r - \lambda = 0. \quad (\text{B15})$$

Notice that the points $\vec{m}_r$ that satisfy this equation define an ellipse, $\mathcal{E}$, with focus at $-A^{-1}\vec{V}$. Notice also that the parameter λ at the maximum is the value of d in (B10) (just multiply (B15) by c_r and sum over r taking into account the constraints $\Lambda = 0$ and $\vec{\Gamma} = 0$).

Finally, consider the variations of $\vec{m}_r$ in (B11). We

obtain

$$c_r \left(A \frac{\vec{V} + A\vec{m}_r}{|\vec{V} + A\vec{m}_r|} - \vec{\gamma} \right) = \omega_r \vec{m}_r, \quad (\text{B16})$$

which means that the vector inside the brackets is proportional to $\vec{m}_r$. Note that condition $\Omega_r = 0$ defines a unit circle, and the orthogonal vector to this curve is $\vec{m}_r$. So we only need to prove that the orthogonal vector at point $\vec{m}_r$ of the ellipse $\mathcal{E}$ defined in (B15) has precisely this direction —this follows straightforwardly taking variations with respect to $\vec{m}_r$ in (B15). Therefore, the solution is given by the tangency points of ellipse $\mathcal{E}$ and circle $\Omega_r = 0$. There are only two such points, they are in opposite directions, and all constraints and maximization equations are satisfied with $c_{1,2} = 1/2$. This proves that the optimal measurements in the greedy scheme are indeed von Neumann's [42]. Notice that this is a stronger statement than it looks: local measurements with a larger number of outcomes will perform worse.

-
- [1] W. K. Wootters and W. H. Zurek, *Nature* **299**, 802 (1982).
 - [2] A. Peres and W. K. Wootters, *Phys. Rev. Lett.* **66**, 1119 (1991).
 - [3] S. Massar and S. Popescu, *Phys. Rev. Lett.* **74**, 1259 (1995).
 - [4] A. S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory* (North Holland, Amsterdam, 1982).
 - [5] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976).
 - [6] M. Paris and J. Reháček (Eds.), *Quantum State Estimation*, *Lectures Notes on Physics* **649** (Springer, Berlin, 2004).
 - [7] S. L. Braunstein and H. J. Kimble, *Phys. Rev. A* **61**, 042302 (2000).
 - [8] R. Derka, V. Bužek and A. K. Ekert, *Phys. Rev. Lett.* **80**, 1571 (1998).
 - [9] J. I. Latorre, P. Pascual, R. Tarrach, *Phys. Rev. Lett.* **81**, 1351 (1998); S. Ibildir and J. Roland, e-print quant-ph/0410237.
 - [10] N. Gisin and S. Popescu, *Phys. Rev. Lett.* **83**, 432 (1999); S. Massar, *Phys. Rev. A* **62**, 040101 (2000).
 - [11] E. Bagan *et al.*, *Phys. Rev. Lett.* **85**, 5230 (2000); *ibid.*, *Phys. Rev. A* **63**, 052309 (2001); E. Bagan, M. Baig and R. Muñoz-Tapia, *Phys. Rev. A* **64**, 022305 (2001).
 - [12] A. Peres and P. F. Scudo, *Phys. Rev. Lett.* **86**, 4160 (2001).
 - [13] A. Acín, E. Jane and G. Vidal, *Phys. Rev. A* **64**, 050302 (2001); E. Jane, Ph.D. Thesis, University of Barcelona, *unpublished*.
 - [14] A. Peres and P. F. Scudo, *Phys. Rev. Lett.* **87**, 167901 (2001); *ibid.*, *J. Modern Optics* **49**, 1235 (2002); N. H. Lindner, A. Peres and D. R. Terno, *Phys. Rev. A* **68**, 042308 (2003).
 - [15] E. Bagan, M. Baig and R. Muñoz-Tapia, *Phys. Rev. Lett.* **87**, 257903 (2001); *ibid.*, *Phys. Rev. A* **69**, 050303 (2004).
 - [16] E. Bagan, M. Baig and R. Muñoz-Tapia, *Phys. Rev. A* **70**, 030301 (2004).
 - [17] S. D. Bartlett, T. Rudolph and R. W. Spekkens, *Phys. Rev. Lett.* **91**, 027901 (2003).
 - [18] K. R. Jones, *Phys. Rev. A* **50**, 3682 (1994).
 - [19] R. D. Gill and S. Massar, *Phys. Rev. A* **61**, 042312 (2000).
 - [20] D. G. Fischer, S. H. Kienle and M. Freyberger, *Phys. Rev. A* **61**, 032306 (2000).
 - [21] Th. Hannemann *et al.*, *Phys. Rev. A* **65**, 050303 (2002).
 - [22] E. Bagan, M. Baig and R. Muñoz-Tapia, *Phys. Rev. Lett.* **89**, 277904 (2002).
 - [23] E. Bagan, M. Baig, R. Muñoz-Tapia, and A. Rodriguez, *Phys. Rev. A* **69**, 010304 (2004).
 - [24] F. Embacher and H. Narnhofer, *Ann. of Phys (N.Y.)* **311**, 220 (2004).
 - [25] M. Hayashi, *A Linear Programming Approach to Attainable Cramér-Rao Type Bound*, in *Quantum Communication, Computing and Measurement*, edited by O. Hirota, A. S. Holevo and C. M. Caves, (Plenum Publishing, New York, 1997); M. Hotta and M. Ozawa, e-print quant-ph/0401187.
 - [26] K. Matsumoto, *J. Phys. A* **35**, 3111 (2002); K. Usami *et al.*, *Phys. Rev. A* **68**, 022314 (2003); M. Hayashi and K. Matsumoto, e-print quant-ph/0308150.
 - [27] C. M. Caves, C. A. Fuchs and P. Rungta, *Found. of Phys. Lett.* **14**, **199** (2001); J. Batle *et al.*, *Optics and Spectroscopy* **94**, 700 (2003).
 - [28] H. M. Wiseman and R. B. Killip, *Phys. Rev. A* **56**, 944 (1997); *ibid.* **57**, 2169 (1998).
 - [29] D. W. Berry, H. M. Wiseman and J. K. Breslin, *Phys. Rev. A* **63**, 053804 (2001); D. W. Berry and H. M. Wiseman, *Phys. Rev. A* **65**, 043803 (2002).

- [30] U. Leonhardt, *Measuring the Quantum State of Light* (Cambridge Univ. Press, Cambridge, England, 1997).
- [31] A. G. White *et al*, Phys. Rev. Lett. **83**, 3103 (1999); D. F. V. James *et al*, Phys. Rev. A **64**, 052313 (2001); R. T. Thew *et al*, Phys. Rev. A **66**, 012303 (2002); J. B. Altepeter *et al*, Phys. Rev. Lett. **90**, 193601 (2003); E. Skovsen, H. Stapelfeldt, S. Juhl and K. Molmer, Phys. Rev. Lett. **91**, 090406 (2003).
- [32] Y. I. Bogdanov *et al*, Phys. Rev. A **70**, 042303 (2004); H. F. Hofmann and S. Takeuchi, Phys. Rev. A **69**, 042108 (2004).
- [33] Z. Hradil, Phys. Rev. A **55**, 1561 (1997); K. Banaszek, Phys. Rev. A **59**, 4797 (1999); Z. Hradil, J. Summhammer and H. Rauch, Phys. Lett. A **261**, 20 (1999); K. Banaszek *et al*, Phys. Rev. A **61**, 010304 (2000); Z. Hradil, *et al*, Phys. Rev. A **62**, 014101 (2000); J. Fiurásek and Z. Hradil, Phys. Rev. A **63**, 020101 (2001).
- [34] A. R. Edmonds, *Angular Momentum in Quantum Mechanics*, (Princeton University Press, 1960).
- [35] A. Monras, Master Thesis (Universitat Autònoma de Barcelona, Barcelona 2004).
- [36] D. Ivanovic, J. Phys. A **14**, 3241 (1981); W. K. Wootters, B. D. Fields, Ann. Phys. (N.Y.) **191**, 363 (1989).
- [37] T. Cover and J. Thomas, *Elements of Information Theory* (Wiley, New York, 1991)
- [38] S. L. Braunstein and C. M. Caves, Phys. Rev. Lett. **72**, 3439 (1994).
- [39] H. Cramer, *Mathematical Methods of Statistics* (Princeton Univ. Press, Princeton, 1946).
- [40] The generalization of this formula to arbitrary N , i.e., $\vec{m}(\chi_N) = (-1)^{i_N} \sum_{k=1}^{N-2} \vec{m}(\chi_k) \times \vec{m}(\chi_{N-1}) / \sqrt{N-2}$, provides a set of points that are roughly isotropically distributed in the sphere, which may be interesting in other contexts.
- [41] Although one can consider more sophisticated estimators, such as MLE or OG, they will not improve significantly the estimation.
- [42] For the 3D case we just have to replace ellipses by ellipsoids and circles by spheres.